

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 453/02, A61K 31/435	A1	(11) International Publication Number: WO 94/11368 (43) International Publication Date: 26 May 1994 (26.05.94)
(21) International Application Number: PCT/US93/06624 (22) International Filing Date: 19 July 1993 (19.07.93) (30) Priority data: 07/975,244 12 November 1992 (12.11.92) US (60) Parent Application or Grant (63) Related by Continuation US 07/975,244 (CON) Filed on 12 November 1992 (12.11.92) (71) Applicant (for all designated States except US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US).		(72) Inventors; and (75) Inventors/Applicants (for US only) : ITO, Fumitaka [JP/JP]; 1-1-2 Sakuragaoka, Taketoy-cho, Chita-gun, Aichi 470-23 (JP). KONDO, Hiroshi [JP/JP]; 2-12 Kariyado-cho, Handa, Aichi 475 (JP). NAKANE, Masami [JP/JP]; 6-28 Kawanahonmachi, Showa-ku, Nagoaya, Aichi 466 (JP). SHIMADA, Kaoru [JP/JP]; 171 Dannoue, Iwadzo-cho, Okazaki, Aichi 444-21 (JP). (74) Agents: RICHARDSON, Peter, C. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US). (81) Designated States: AU, CA, JP, KR, NO, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: QUINUCLIDINE DERIVATIVE AS SUBSTANCE P ANTAGONIST (57) Abstract This invention relates to the quinuclidine derivative (2S, 3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine and its pharmaceutically acceptable salts. These compounds are substance P antagonists and are useful in the treatment of gastrointestinal disorders, inflammatory disorders, central nervous system disorders and pain.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

-1-

5

QUINUCLIDINE DERIVATIVE AS SUBSTANCE P ANTAGONIST

Background of the Invention

The present invention relates to the novel quinuclidine derivatives (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-
10 diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine, pharmaceutical compositions comprising such compound and the use of such compound in the treatment and prevention of inflammatory and central nervous system disorders, as well as several other disorders. The pharmaceutically active
15 compound of this invention is a substance P receptor antagonists.

Substance P is a naturally occurring undecapeptide belonging to the tachykinin family of peptides, the latter being named because of their prompt stimulatory action on
20 smooth muscle tissue. More specifically, substance P is a pharmacologically active neuropeptide that is produced in mammals (having originally been isolated from gut) and possesses a characteristic amino acid sequence that is illustrated by D. F. Veber et al. in U.S. Patent No.
25 4,680,283. The wide involvement of substance P and other tachykinins in the pathophysiology of numerous diseases has been amply demonstrated in the art. For instance, substance P has recently been shown to be involved in the transmission of pain or migraine (see B.E.B. Sandberg et al., Journal of
30 Medicinal Chemistry, 25, 1009 (1982)), as well as in central nervous system disorders such as anxiety and schizophrenia, in respiratory and inflammatory diseases such as asthma and rheumatoid arthritis, respectively, in rheumatic diseases such as fibrositis, and in gastrointestinal disorders and
35 diseases of the GI tract such as ulcerative colitis and Crohn's disease, etc. (see D. Regoli in "Trends in Cluster Headache," edited by F. Sicuteri et al., Elsevier Scientific Publishers, Amsterdam, pp. 85-95 (1987)).

The quinuclidine derivative of this invention is
40 referred to generically in United States Patent 5,162,339, which issued to John A. Lowe III on November 11, 1992.


-2-

Quinuclidine, piperidine, and azanorbornane derivatives and related compounds that exhibit activity as substance P receptor antagonists are referred to in United States Patent Application 724,268, filed July 1, 1991, PCT Patent Application PCT/US 91/02853, filed April 25, 1991, PCT Patent Application PCT/US 91/03369, filed May 14, 1991, PCT Patent Application PCT/US 91/05776, filed August 20, 1991, PCT Patent Application PCT/US 92/00113, filed January 17, 1992, PCT Patent Application PCT/US 92/03571, filed May 5, 1992, PCT Patent Application PCT/US 92/03317, filed April 28, 1992, PCT Patent Application PCT/US 92/04697, filed June 11, 1992, United States Patent Application 766,488, filed September 26, 1991, United States Patent Application 790,934, filed November 12, 1991, PCT Patent Application PCT/US 92/04002, filed May 19, 1992, Japanese Patent Application No. 065337/92, filed March 23, 1992, and United States Patent Application 932,392, filed August 19, 1992.

Summary of the Invention

The present invention relates to the quinuclidine derivative (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine and its pharmaceutically acceptable salts.

(2S,3S)-N-(5-n-Propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine has the following chemical structure



The present invention also relates to a pharmaceutical composition for treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g.,

-3-

arthritis, psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, hypertension, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis in a mammal, including a human, comprising an amount of (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such condition, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, hypertension, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic neuropathy and multiple sclerosis, disorders related to immune enhancement or

-4-

suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis in a mammal, including a human, comprising administering to said mammal an amount of (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such condition.

The present invention also relates to a pharmaceutical composition for antagonizing the effects of substance P in a mammal, including a human, comprising a substance P antagonizing amount of (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of antagonizing the effects of substance P in a mammal, including a human, comprising administering to said mammal a substance P antagonizing amount of (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine, or a pharmaceutically acceptable salt thereof.

The present invention also relates to a pharmaceutical composition for treating or preventing a disorder in a mammal, including a human, resulting from an excess of substance P, comprising a substance P antagonizing amount of (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a disorder in a mammal, including a human, resulting from an excess of substance P, comprising administering to said mammal a substance P antagonizing amount of (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine, or a pharmaceutically acceptable salt thereof.

-5-

The present invention also relates to a pharmaceutical composition for treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, hypertension, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fasciitis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis in a mammal, including a human, comprising an amount of (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine, or a pharmaceutically acceptable salt thereof, effective in antagonizing the effect of substance P at its receptor site, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, hypertension, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fasciitis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia,

-6-

neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and
5 rheumatic diseases such as fibrositis in a mammal, including a human, comprising administering to said mammal an amount of (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine, or a pharmaceutically acceptable salt thereof, effective in
10 antagonizing the effect of substance P at its receptor site.

The present invention also relates to a pharmaceutical composition for treating or preventing a disorder in a mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance
15 P mediated neurotransmission, comprising an amount of (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine, or a pharmaceutically acceptable salt thereof, effective in antagonizing the effect of substance P at its receptor site,
20 and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a disorder in mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated
25 neurotransmission, comprising administering to said mammal an amount of (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine, or a pharmaceutically acceptable salt thereof, effective in antagonizing the effect of substance P at its receptor site.

30 The present invention also relates to a pharmaceutical composition for treating or preventing a disorder in a mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising an amount of
35 (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine, or a pharmaceutically acceptable salt thereof, effective in

-7-


treating or preventing such disorder, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a disorder in mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising administering to said mammal an amount of (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such disorder.

-8-


Detailed Description of the Invention

(2S,3S)-N-(5-n-Propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine
(hereinafter also referred to as "the active compound of
5 this invention") may be prepared by subjecting a compound of
the formula




II

having the same absolute stereochemistry as the active
compound of this invention, to hydrolytic removal of the
methoxybenzyl group to produce the corresponding compound of
the formula



III

having the same stereochemistry, and then reacting the
compound of formula III so formed with an aldehyde of the
30 formula



IV

-9-

in the presence of a reducing agent.

Hydrolytic removal of the methoxybenzyl group is generally carried out using a strong mineral acid such as hydrochloric, hydrobromic or hydroiodic acid, at a temperature from about room temperature to about the reflux temperature of the acid. Preferably, the reaction is conducted in hydrobromic acid at the reflux temperature. This reaction is usually carried out for a period of about 2 hours.

Alternatively, the hydrolytic removal of the methoxybenzyl group in the above procedure may be replaced by hydrogenolytic removal of such group. Hydrogenolytic removal is generally accomplished using hydrogen in the presence of a metal containing catalyst such as platinum or palladium. This reaction is usually conducted in a reaction inert solvent such as acetic acid or a lower alcohol, at a temperature from about 0°C to about 50°C. The methoxybenzyl group may also be removed, alternatively, by treating the compound of formula II with a dissolving metal such as lithium or sodium in ammonia at a temperature from about -30°C to about 78°C, or with a formate salt in the presence of palladium or with cyclohexane in the presence of palladium.

Preferably, the methoxybenzyl group is removed by treating the compound of formula II with hydrogen in the presence of palladium hydroxide on carbon in methanol containing hydrochloric acid at a temperature of about 25°C.


The resulting compound of formula III may be converted into the active compound of this invention by reaction with the aldehyde of formula IV in the presence of a reducing agent. The reaction is typically carried out using a reducing agent such as sodium cyanoborohydride, sodium triacetoxyborohydride, sodium borohydride, hydrogen and a metal catalyst, zinc and hydrochloric acid, borane dimethylsulfide or formic acid at a temperature from about -60°C to about 50°C. Suitable reaction inert solvents for this reaction include lower alcohols (e.g., methanol,

-10-

ethanol and isopropanol), acetic acid, methylene chloride and tetrahydrofuran (THF). Preferably, the solvent is methylene chloride, the temperature is about 25°C, and the reducing agent is sodium triacetoxymethylborohydride.

5 Alternatively, the reaction of the compound of the formula III with the compound of the formula IV may be carried out in the presence of a drying agent or using an apparatus designed to remove azeotropically the water generated, to produce an imine of the formula

10




15

which is then reacted with a reducing agent as described above, preferably with sodium triacetoxymethylborohydride at about room temperature. The preparation of the imine is generally carried out in a reaction inert solvent such as benzene, xylene or toluene, preferably toluene, at a temperature from about 25°C to about 110°C, preferably at about the reflux temperature of the solvent. Suitable drying agents/solvent systems include titanium tetrachloride/dichloromethane, titanium isopropoxide/dichloromethane and molecular sieves/THF. Titanium tetrachloride/dichloromethane is preferred.

30 The active compound of this invention may also be prepared from a compound of the formula III having the same stereochemistry by reacting the compound of formula III with a compound of the formula


35

-11-



5
10 wherein L is a suitable leaving group (e.g., chloro, bromo, iodo or mesylate). This reaction is generally carried out in a reaction inert solvent such as dichloromethane or THF, preferably dichloromethane, at a temperature from about 0°C to about 60°C , preferably at about 25°C .

15 The active compound of this invention may also be prepared from a compound of the formula III having the same stereochemistry by reacting the compound of formula III with a compound of the formula



20
25
30 wherein L is defined as above or is imidazole, and then reducing the resulting amide. This reaction is typically carried out in an inert solvent such as THF or dichloromethane at a temperature from about -20°C to about 60°C , preferably in dichloromethane at about 0°C . Reduction of the resulting amide is accomplished by treatment with a reducing agent such as borane dimethylsulfide complex, lithium aluminum hydride or diisobutylaluminum hydride in an
35 inert solvent such as ethyl ether or THF. The reaction temperature may range from about 0°C to about the reflux temperature of the solvent. Preferably, the reduction is

-12-

accomplished using borane dimethylsulfide complex in THF at about 60°C.

The active compound of this invention and its pharmaceutically acceptable salts are useful as substance P antagonists, i.e., they possess the ability to antagonize the effects of substance P at its receptor site in mammals, and therefore they are able to function as therapeutic agents in the treatment of the aforementioned disorders and diseases in an afflicted mammal.

10 (2S,3S)-N-(5-n-Propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine is basic in nature and therefore capable of forming a wide variety of different salts with various inorganic and organic acids. Although such salts must be pharmaceutically acceptable for
15 administration to animals, it is often desirable in practice to initially isolate the active compound of this invention from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent and
20 subsequently convert the latter free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the active compound of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the chosen mineral or
25 organic acid in an aqueous solvent medium or in a suitable organic solvent, such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is readily obtained.

The active compound of this invention and its
30 pharmaceutically acceptable salts exhibit substance P receptor binding activity and therefore are of value in the treatment and prevention of a wide variety of clinical conditions the treatment or prevention of which are effected or facilitated by a decrease in substance P mediated
35 neurotransmission. Such conditions include inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), anxiety, depression or

-13-

dysthymic disorders, colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, hypertension, vasospastic diseases such as angina, migraine
5 and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fasciitis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia,
10 neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis. Hence, these
15 compounds are readily adapted to therapeutic use as substance P antagonists for the control and/or treatment of any of the aforesaid clinical conditions in mammals, including humans.

The active compound of this invention and its
20 pharmaceutically acceptable salts can be administered via either the oral, parenteral or topical routes. In general, these compounds are most desirably administered in dosages ranging from about 0.5 mg to about 500 mg per day, although variations will necessarily occur depending upon the weight
25 and condition of the subject being treated and the particular route of administration chosen. Variations may occur depending upon the species of animal being treated and its individual response to said medicament, as well as on the type of pharmaceutical formulation chosen and the time
30 period and interval at which such administration is carried out. In some instances, dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect, provided that such larger
35 doses are first divided into several small doses for administration throughout the day.

-14-

The active compound of this invention and its pharmaceutically acceptable salts may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by either of the three routes previously indicated, and such administration may be carried out in single or multiple doses. More particularly, such compounds can be administered in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, lozenges, troches, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various non-toxic organic solvents, etc. Moreover, oral pharmaceutical compositions can be suitably sweetened and/or flavored. In general, the active compound of this invention or a pharmaceutically acceptable salt thereof is present in such dosage forms at concentration levels ranging from about 5.0% to about 70% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine may be employed along with various disintegrants such as starch (and preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or

-15-

dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

5 For parenteral administration, solutions of the active compound of this invention, or a pharmaceutically acceptable salt thereof, in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered (preferably pH greater than 8)
10 if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under
15 sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art.

Additionally, it is also possible to administer the active compound of this invention and its pharmaceutically
20 acceptable salts topically when treating inflammatory conditions of the skin and this may preferably be done by way of creams, jellies, gels, pastes, ointments and the like, in accordance with standard pharmaceutical practice.

The activity of the active compound of this invention
25 and its pharmaceutically acceptable salts as substance P receptor antagonists may be determined by its ability to inhibit the binding of substance P at its receptor sites in bovine caudate tissue, employing radioactive ligands to visualize the tachykinin receptors by means of
30 autoradiography. The substance P antagonizing activity of such compounds may be evaluated by using the standard assay procedure described by M. A. Cascieri et al., as reported in the Journal of Biological Chemistry, Vol. 258, p. 5158 (1983). This method essentially involves determining the
35 concentration of the active compound of this invention, or a pharmaceutically acceptable salt thereof, required to reduce by 50% the amount of radiolabelled substance P

-16-

ligands at their receptor sites in said isolated cow tissues, thereby affording characteristic IC_{50} values for the compound tested.

In this procedure, bovine caudate tissue is removed from a $-70^{\circ}C$ freezer and homogenized in 50 volumes (w./v.) of an ice-cold 50 mM Tris (i.e., trimethamine which is 2-amino-2-hydroxymethyl-1,3-propanediol) hydrochloride buffer having a pH of 7.7. The homogenate is centrifuged at 30,000 x G for a period of 20 minutes. The pellet is resuspended in 50 volumes of Tris buffer, rehomogenized and then recentrifuged at 30,000 x G for another twenty-minute period. The pellet is then resuspended in 40 volumes of ice-cold 50 mM Tris buffer (pH 7.7) containing 2 mM of calcium chloride, 2 mM of magnesium chloride, 40 g/ml of bacitracin, 4 μ g/ml of leupeptin, 2 μ g of chymostatin and 200 g/ml of bovine serum albumin. This step completes the production of the tissue preparation.

The radioligand binding procedure is then carried out in the following manner, viz., by initiating the reaction via the addition of 100 μ l of the test compound made up to a concentration of 1 μ M, followed by the addition of 100 μ l of radioactive ligand made up to a final concentration 0.5 mM and then finally by the addition of 800 μ l of the tissue preparation produced as described above. The final volume is thus 1.0 ml, and the reaction mixture is next vortexed and incubated at room temperature (ca. $20^{\circ}C$) for a period of 20 minutes. The tubes are then filtered using a cell harvester, and the glass fiber filters (Whatman GF/B) are washed four times with 50 mM of Tris buffer (pH 7.7), with the filters having previously been presoaked for a period of two hours prior to the filtering procedure. Radioactivity is then determined in a Beta counter at 53% counting efficiency, and the IC_{50} values are calculated by using standard statistical methods.

The anti-psychotic activity of the active compound of this invention and its pharmaceutically acceptable salts as neuroleptic agents for the control of various psychotic

-17-

disorders may be determined primarily by a study of its ability to suppress substance P-induced or substance P agonist induced hypermotility in guinea pigs. This study is carried out by first dosing the guinea pigs with a control compound or with an appropriate test compound of the present invention, then injecting the guinea pigs with substance P or a substance P agonist by intracerebral administration via canula and thereafter measuring their individual locomotor response to said stimulus.

10 The present invention is illustrated by the following example. It will be understood, however, that the invention is not limited to the specific details of this example.

EXAMPLE

15 (2S,3S)-N-(2-Methoxy-5-n-propylphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine methanesulfonate

To a solution of a 2-methoxy-5-n-propylbenzaldehyde (370 mg, 2.06 mmol) (prepared by Duff's formylation of 4-n-propylanisole, as described in Synth. Common., 15, 61 (1985)) and (2S,3S)-diphenylmethyl-1-azabicyclo-[2,2,2]octan-3-amine (1.71 mmol) in methylene chloride (20 ml) was added in portions sodium triacetoxyborohydride (510 mg, 2.39 mmol). The mixture was stirred until the amine disappeared. The solution was carefully neutralized with an ice cooled 2N sodium hydroxide solution. The organic layer was washed with water, dried over magnesium sulfate, and concentrated to give the product (840 mg), which was chromatographed on a silica gel column. ¹H NMR (270 MHz, CDCl₃, ppm): δ 10.45 (s, 1H), 7.64 (d, J=2.6 Hz, 1H), 7.37 (dd, J=8.4, 2.6 Hz, 1H), 6.91 (d, J=8.4 Hz, 1H), 3.91 (s, 3H), 2.56 (t, J=7.3 Hz, 2H), 1.62 (m, 2H), 0.92 (t, J=7.3 Hz, 3H).

Methanesulfonic acid (96 μl) was added to the product (650 mg). The precipitate was recrystallized from acetone to give the analytical pure product (240 mg).

35 M.P.: 237-241°C (acetone).

Analysis calc'd for C₃₁H₃₈N₂O•CH₃SO₃H•1/3H₂O: C, 69.04%; H, 7.72%; N, 5.03%. Found C, 68.96%; H, 7.88%; N, 4.99%.

-18-

¹H NMR (270 MHz, CDCl₃, ppm): δ 7.36-7.04 (m, 10H), 6.94 (dd, J=8.4, 2.5 Hz, 1H), 6.62 (d, J=8.4 Hz, 1H), 6.46 (d, J=2.5 Hz, 1H), 4.50 (d, J=12.0 Hz, 1H), 3.68 (dd, J=12.0, 8.0 Hz, 1H), 3.58 (d, J=14.0 Hz, 1H), 3.54 (s, 3H), 3.22 (d, J=14.0 Hz, 1H), 3.27-3.15 (m, 1H), 2.92 (dd, J=8.0, 4.0 Hz, 1H), 2.76 (m, 2H), 2.60 (m, 1H), 2.44 (t, J=7.4 Hz, 2H), 2.07 (m, 1H), 1.97-1.89 (m, 1H), 1.70-1.45 (m, 4H), 1.31-1.20 (m, 1H), 0.94 (t, J=7.4 Hz, 3H).

¹³C NMR (67.5 MHz, CDCl₃, ppm): δ 155.5, 145.6, 143.3, 10 134.1, 129.4, 128.9, 128.3, 127.6, 127.5, 127.4, 126.3, 125.8, 109.8, 61.9, 55.3, 54.6, 49.4, 49.2, 46.5, 41.9, 37.1, 25.5, 24.9, 24.7, 19.9, 13.9.

IR (KBr): 3,410 (br.), 1,502 (s), 1,455 (m), 1,300-1,100 (br, m), 753 (s), 710(s).

-19-

CLAIMS

1. (2S,3S)-N-(5-n-propyl-2-methoxyphenyl)methyl-2-diphenylmethyl-1-azabicyclo[2.2.2]octan-3-amine and its pharmaceutically acceptable salts.

5 2. A pharmaceutical composition for treating or preventing a condition selected from the group consisting of inflammatory diseases, anxiety, colitis, depression or dysthymic disorders, psychosis, pain, allergies, chronic obstructive airways disease, hypersensitivity disorders,
10 hypertension, vasospastic diseases, fibrosing and collagen diseases, reflex sympathetic dystrophy, addiction disorders, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders, disorders related to immune enhancement or suppression and rheumatic diseases in
15 a mammal, comprising an amount of a compound according to claims 1 that is effective in preventing or treating such condition and a pharmaceutically acceptable carrier.

3. A method of treating or preventing a condition selected from the group consisting of inflammatory diseases
20 anxiety, colitis, depression or dysthymic disorders, psychosis, pain, allergies, chronic obstructive airways disease, hypersensitivity disorders, hypertension, vasospastic diseases, fibrosing and collagen diseases, reflex sympathetic dystrophy, addiction disorders, stress
25 related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders, disorders related to immune enhancement or suppression and rheumatic diseases in a mammal, comprising administering to a mammal in need of such treatment or prevention an amount of a compound according to
30 claim 1 that is effective in preventing or treating such condition.

4. A pharmaceutical composition for antagonizing the effects of substance P in a mammal, comprising a substance P antagonizing effective amount of a compound according to
35 claim 1 and a pharmaceutically acceptable carrier.

5. A method of antagonizing the effects of substance P in a mammal, comprising administering to said mammal a

-20-

substance P antagonizing effective amount of a compound according to claim 1.

6. A pharmaceutical composition for treating or preventing a condition in a mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising an amount of a compound according to claim 1 that is effective in antagonizing the effect of substance P at its receptor site and a pharmaceutically acceptable carrier.

7. A method of treating or preventing a condition in a mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising administering to a mammal in need of such treatment or prevention an amount of a compound according to claim 1 that is effective in antagonizing the effect of substance P at its receptor site.

8. A pharmaceutical composition for treating or preventing a condition in a mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising an amount of a compound according to claim 1 that is effective in treating or preventing such condition and a pharmaceutically acceptable carrier.

9. A method of treating or preventing a condition in a mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising administering to a mammal in need of such treatment or prevention an amount of a compound according to claim 1 that is effective in treating or preventing such condition.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 93/06624

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C07D453/02; A61K31/435		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification Systems	Classification Symbols	
Int.Cl. 5	C07D	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X	WO,A,9 005 729 (PFIZER INC.) 31 May 1990 cited in the application * example 4 *	1,2,4
A	----- TRENDS IN PHARMACOLOGICAL SCIENCES vol. 13, July 1992, pages 266 - 269 K.J. WATLING 'Nonpeptide antagonists herald a new era in tachykinin research' * page 266, compound CP 96345 *	1,2,4
P,X	----- WO,A,9 221 677 (PFIZER INC.) 10 December 1992 * complete document *	1,2,4
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
02 SEPTEMBER 1993		14. 09. 93
International Searching Authority		Signature of Authorized Officer
EUROPEAN PATENT OFFICE		VAN BIJLEN H.

INTERNATIONAL SEARCH REPORT

national application No.

PCT/US 93/06624

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 3,5,7 and 9 are directed to a method of treatment of (diagnostic method practised on) the human/animal body, the search has been carried out and based on the alleged effects of the compound/ composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

US 9306624
SA 76697

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

02/09/93

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9005729	31-05-90	WO-A- 9005525	31-05-90
		CA-A- 2003441	23-05-90
		EP-A- 0409931	30-01-91
		US-A- 5162339	10-11-92

WO-A-9221677	10-12-92	AU-A- 1990192	08-01-93
		CN-A- 1067428	30-12-92
